

# Pituitary Adenylate Cyclase-Activating Polypeptide Induces the Voltage-Independent Activation of Inward Membrane Currents and Elevation of Intracellular Calcium in HIT-T15 Insulinoma Cells\*

COLIN A. LEECH, GEORGE G. HOLZ, AND JOEL F. HABENER†

Laboratory of Molecular Endocrinology, Howard Hughes Medical Institute, and the Diabetes Unit (G.G.H.), Harvard Medical School, Massachusetts General Hospital, Boston, Massachusetts 02114

## ABSTRACT

The secretion of insulin by pancreatic  $\beta$ -cells is controlled by synergistic interactions of glucose and hormones of the glucagon-related peptide family, of which pituitary adenylate cyclase-activating polypeptide (PACAP) is a member. Here we show by simultaneous recording of intracellular calcium ion ( $[Ca^{2+}]_i$ ) and membrane potential that both PACAP-27 and PACAP-38 depolarize HIT-T15 cells and raise  $[Ca^{2+}]_i$ . PACAP stimulation can result in membrane depolarization by two distinct mechanisms: 1) PACAP reduces the membrane conductance and increases membrane excitability; and 2) PACAP activates a pronounced inward current that is predominantly a  $Na^+$  current, blockable by  $La^{3+}$ , and which exhibits a reversal potential of about  $-28$  mV. Activation of this current does not require membrane

depolarization, because the response is observed when cells are held under voltage clamp at  $-70$  mV. This current may result from the cAMP-dependent activation of nonspecific cation channels because the current is also observed in response to forskolin or membrane-permeant analogs of cAMP.

We also suggest that PACAP raises  $[Ca^{2+}]_i$  and stimulates insulin secretion by three distinct mechanisms: 1) depolarization activates  $Ca^{2+}$  influx through L-type voltage-dependent calcium channels, 2) mobilization of intracellular  $Ca^{2+}$  stores, and 3) entry of  $Ca^{2+}$  via voltage-independent  $Ca^{2+}$  channels. These effects of PACAP may play an important role in a neuro-entero-endocrine loop regulating insulin secretion from pancreatic  $\beta$ -cells during the transition period from fasting to feeding. (*Endocrinology* 136: 1530–1536, 1995)

PITUITARY adenylate cyclase-activating polypeptide (PACAP) is a neuropeptide structurally related to the peptide family consisting of glucagon, secretin, vasoactive intestinal peptide, and others (1). Two isoforms of PACAP, PACAP-27 and PACAP-38, which share the same 27 N-terminal amino acids, are derived by tissue-specific posttranslational processing of a 176-amino acid precursor (2). In the endocrine pancreas, PACAP is produced and secreted from nerve endings that innervate the islets of Langerhans (3), and PACAP is an extremely potent stimulator of glucose-dependent insulin secretion (4). Multiple isoforms of the PACAP receptor (PACAP-R) have been isolated and sequenced, including, in insulin-secreting  $\beta$ -cells, the type 3 PACAP-R (PACAP-R3), which is homologous to the rat vasoactive intestinal peptide<sub>2</sub> receptor (5, 6). The type 1 PACAP-R has been shown to stimulate both cAMP accumulation ( $EC_{50}$  3 nM) and [<sup>3</sup>H]inositol phosphate production ( $EC_{50}$ , 20 nM) in the gonadotrope-derived  $\alpha$ T3-1 cell line (7). Similarly, when expressed in *Xenopus* oocytes, PACAP-R3 can couple to phospholipase-C as well as to adenylyl cyclase (5), but such dual coupling of PACAP receptors has yet to be demonstrated in insulinoma cells or pancreatic  $\beta$ -cells. However, the observation that nitrendipine blocks PACAP-induced

elevation of intracellular calcium ion ( $[Ca^{2+}]_i$ ) and insulin secretion (4) suggests that mobilization of intracellular  $Ca^{2+}$  stores through activation of phospholipase-C is not the primary signaling pathway of PACAP in  $\beta$ -cells and that stimulation of cAMP production may play a more dominant role.

In perfused rat pancreas, the two isoforms of PACAP, PACAP-27 and PACAP-38, induce insulin secretion in a glucose-dependent manner. PACAP-38 is more potent than PACAP-27 at 5.5 mM extracellular glucose, but the two forms are equally potent at 8.3 mM glucose (8). In rat  $\beta$ -cells, both PACAP-27 and PACAP-38 potentiate glucose-dependent insulin secretion at concentrations as low as  $10^{-14}$ – $10^{-13}$  M (4). These actions of PACAP are reported to be mediated through an increase in the activity of L-type voltage-dependent calcium channels (VDCCs), because nitrendipine, a specific blocker of L-type channels, abolishes the stimulatory effects of PACAP on both  $[Ca^{2+}]_i$  and insulin secretion. However, the effects of PACAP on membrane currents or membrane potential were not examined (4).

As PACAP stimulates cAMP production, it is of interest to determine what effect it might exert on ion channels that are known to be regulated by this cyclic nucleotide, either through direct effects of cAMP on channel gating or via effects mediated by cAMP-dependent protein phosphorylation. Recent studies suggest that cAMP (and  $Ca^{2+}$ ) activate a nonspecific cation channel (Ca-NS channels) in the rat insulinoma cell line, CRI-G1 (9). The activation of this current in isolated inside-out membrane patches does not appear to be mediated by protein kinase-A (PKA) and requires non-

Received September 28, 1994.

Address all correspondence and requests for reprints to: Dr. Colin A. Leech, Laboratory of Molecular Endocrinology, Wellman 320, Massachusetts General Hospital, Boston, Massachusetts 02114.

\* This work was supported by NIH Grants DK-30834, DK-30457 (to J.F.H.), and DK-45817 (to G.G.H.).

† Investigator, Howard Hughes Medical Institute.

physiological cytoplasmic  $[Ca^{2+}]_i$  ( $>0.1$  mM) (9, 10) (the  $[Ca^{2+}]_i$  required in the intact cell may be much lower). The activation of this current *in vivo* would be expected to cause membrane depolarization and a rise in  $[Ca^{2+}]_i$ , thereby triggering insulin secretion.

Glucagon-like peptide-1 (GLP-1) is a potent incretin hormone produced in and released from entero-endocrine cells in the small intestine in response to the ingestion of nutrients. The stimulation of insulin secretion by GLP-1 appears to be mediated by increases in cAMP levels, which potentiate glucose-induced closure of ATP-sensitive K-channels (ATP-K channels) (11). This closure leads to membrane depolarization and activation of VDCCs, resulting in a rise of  $[Ca^{2+}]_i$ , which triggers insulin secretion. GLP-1 is a member of the group of peptides related in structure to PACAP, and the glucose-dependent insulinotropic actions of PACAP might be expected to result from a similar mechanism of action. In addition to these effects, cAMP has been reported to enhance the current through L-type  $Ca^{2+}$  channels, by slowing their inactivation, and to have direct stimulatory effects on exocytosis (12).

The aim of the present study was to examine further the signal transduction pathway of PACAP in HIT-T15 cells using fura-2 fluorescence ratio measurements of  $[Ca^{2+}]_i$  combined with perforated patch recording (13) of the membrane potential or membrane current. We show that PACAP induces depolarization of HIT-T15 cells by reducing the membrane conductance, raising the excitability of cells, and voltage-independent activation of an inward current that is predominantly carried by  $Na^+$ . Depolarization of the cell will activate VDCCs and produce a rise in  $[Ca^{2+}]_i$ . However, PACAP also raises  $[Ca^{2+}]_i$  in cells held under voltage clamp, negative to the threshold for activation of VDCCs, by mobilization of intracellular stores and activation of a  $Ca^{2+}$  release-activated  $Ca^{2+}$ -current ( $I_{CRAC}$ ).

As PACAP is reported to be the most potent insulinotropic peptide known (4), we propose that PACAP may play an important role in a neuro-entero-endocrine loop to potentiate insulin secretion during the fasting to feeding transition, before intestinal incretins, such as GLP-1, have an effect.

## Materials and Methods

HIT-T15 cells (passages 64–78), obtained from the American Type Culture Collection (Rockville, MD), were cultured in F-12 medium containing 10 mM glucose and supplemented with 10% horse serum, 2.5% fetal bovine serum, 100 U/ml penicillin, and 100  $\mu$ g/ml streptomycin. Cells were transferred onto glass coverslips coated with 1 mg/ml of Concanavalin-A (type V, Sigma Chemical Co., St. Louis, MO), which facilitates adherence of the cells to the glass. Experiments were performed 2–4 days postplating.

### Measurement of $[Ca^{2+}]_i$

HIT-T15 cells were prepared for measurement of  $[Ca^{2+}]_i$  by washing three times with PBS to remove culture medium and then placed in 5 ml extracellular recording solution (see below) containing 0.8 mM glucose, 2% fetal bovine serum, 0.03% Pluronic F-127, and 5  $\mu$ M fura-2/AM (Molecular Probes, Eugene, OR). Cells were loaded in this solution for 1.5–2 h at room temperature (20–22 C). Loading HIT-T15 cells at room temperature, rather than at 37 C, substantially reduces the degree of intracellular compartmentalization of fura-2 (14). Glass coverslips with fura-2-loaded adherent cells formed the base of a recording chamber

mounted on a Peltier temperature-controlled (32 C) stage of a Zeiss IM35 microscope (Zeiss, Thornwood, NY) equipped with a dual excitation video imaging system (IonOptix Corp., Milton, MA).  $[Ca^{2+}]_i$  was estimated at 1-sec intervals from the ratio of 510-nm emission fluorescence resulting from excitation by 350- and 380-nm wavelength light from the following equation (15):  $[Ca^{2+}]_i = K_d\beta(R - R_{min})/(R_{max} - R)$ , where  $K_d$  is the dissociation constant of fura-2 (224 nM),  $\beta$  is the ratio of 380 nm induced fluorescence of free/bound fura-2,  $R$  is the measured ratio of 350 nm/380 nm fluorescence, and  $R_{min}$  and  $R_{max}$  are 350 nm/380 nm fluorescence ratios in zero  $[Ca^{2+}]_i$  and saturating  $Ca^{2+}$  concentrations, respectively. Values for  $\beta$ ,  $R_{min}$ , and  $R_{max}$  were obtained by *in vitro* calibrations using penta-potassium fura-2.

### Electrophysiology


Cells were bathed in extracellular recording solution containing 138 mM NaCl, 5.6 mM KCl, 2.6 mM  $CaCl_2$ , 1.2 mM  $MgCl_2$ , and 10 mM HEPES-NaOH (pH 7.4). D-Glucose was added to this solution, as specified in the text. In some experiments,  $Na^+$ -free solutions were produced by substituting NaCl with *N*-methyl-D-glucamine (NMG), and the pH was adjusted to 7.4 using HCl. Nominally  $Ca^{2+}$ -free solutions were produced by substituting  $MgCl_2$  for  $CaCl_2$ . The resting potential of cells was measured under current clamp, or the cells were held under voltage clamp, using the perforated patch (13, 16) configuration. Patch pipettes pulled from Kimax-51 glass capillaries (Kimble Glass Inc., Vineland, NJ) were fire polished and tip-dipped in a solution of 95 mM  $K_2SO_4$ , 7 mM  $MgCl_2$ , and 5 mM HEPES-NaOH (pH 7.4), then back-filled with the same solution containing nystatin (240  $\mu$ g/ml). The pipette was connected to an EPC-9 (Heka Electronics, Lambrecht, Germany) patch clamp amplifier, and the access resistance and cell capacitance were monitored after seal formation. The series resistance was compensated for by 60–80%. The pipette solution did not contain fura-2 and was nominally  $Ca^{2+}$  free. A sudden rise in  $[Ca^{2+}]_i$  and a simultaneous decrease in fluorescence intensity provided a useful marker for accidental rupture of the cell membrane, whereupon the experiment was terminated. Test solutions were applied by pneumatic pressure ejection from micropipettes using a General Valves Picospritzer II (Fairfield, NJ). Forskolin, 8-bromo-cAMP (8Br-cAMP), and PACAP-27 and -38 were obtained from Sigma Chemical Co. (St. Louis, MO). Sp-cAMPs was obtained from BioLog (La Jolla, CA).

## Results

Both PACAP-27 (Fig. 1, A and B) and PACAP-38 (Fig. 1C) induced rises in  $[Ca^{2+}]_i$  of HIT-T15 cells preequilibrated in a fixed concentration (2 mM) of glucose, consistent with the presence of PACAP-R3 receptors in these cells (5). Responses to PACAP were of variable duration and were either transient (Fig. 1A) or prolonged (Fig. 1, B and C). Similar responses were seen after application of forskolin (100  $\mu$ M), 8Br-cAMP (1 mM), or Sp-cAMPs (100  $\mu$ M; Fig. 1D). By analogy with the effects of the related peptide, GLP-1, on rat  $\beta$ -cells (11), PACAP might be expected to increase  $[Ca^{2+}]_i$  by potentiating the glucose-dependent closure of ATP-K channels, resulting in membrane depolarization and opening of VDCCs. The membrane conductance of a HIT-T15 cell was monitored in perforated patch voltage clamp by holding the cell at  $-80$  mV and applying 1-sec steps of  $\pm 20$  mV. In this experiment, 10 nM PACAP-27 produced about a 50% decrease in membrane conductance, with only a small effect on the holding current ( $I_h$ ). This observation suggests that glucose-dependent inhibition of the ATP-sensitive K current by PACAP contributes to its ability to generate membrane depolarization (see below) and a consequent rise in  $[Ca^{2+}]_i$ , as previously shown for GLP-1 (11).

Simultaneous recordings of membrane potential and

FIG. 1. Both PACAP-27 and PACAP-38 induce rises in  $[Ca^{2+}]_i$  in HIT cells. A and B, The rise in  $[Ca^{2+}]_i$  in HIT cells after application of 2 nM PACAP-27 for 10 sec. C, A record from a HIT cell that responded to the application of 10 nM PACAP-38 for 10 sec. These responses illustrate the variability of the time course of the effect of changes in  $[Ca^{2+}]_i$  on PACAP stimulation, which may be transient (A) or prolonged (B and C). Similar changes in  $[Ca^{2+}]_i$  could be elicited by the application of membrane-permeant cAMP analogs, such as Sp-cAMPs (100  $\mu$ M; D). The cells were bathed in 2 mM glucose saline.



$[Ca^{2+}]_i$  in current-clamped HIT-T15 cells show that PACAP-27 causes membrane depolarization, the triggering of action potentials, and a simultaneous rise in  $[Ca^{2+}]_i$  (Fig. 3, A and B). However, the same cell held under voltage clamp at  $-70$  mV, negative to the threshold for activation of VDCCs (17), responded to a second application of PACAP-27 with a pronounced inward current (Fig. 3C) and a much larger and more prolonged rise in  $[Ca^{2+}]_i$  (Fig. 3D). This type of response was observed more frequently than that illustrated in Fig. 2. The larger rise in  $[Ca^{2+}]_i$  observed under voltage clamp, compared to current clamp, in this cell might be explained by the increased electrical driving force for cation entry through open channels, because under voltage clamp recording conditions, the membrane potential was held at  $-70$  mV, whereas under current clamp recording conditions, the membrane potential shifts to about  $-20$  mV.

Removal of extracellular  $Ca^{2+}$  ( $Ca^{2+}_o$ ) during the response to PACAP reversibly abolished the rise in  $[Ca^{2+}]_i$  (Fig. 4A).





FIG. 2. PACAP causes an initial reduction in membrane conductance. A HIT cell was held under perforated patch voltage clamp at  $-80$  mV, and 1-sec steps of  $\pm 20$  mV were applied to monitor membrane conductance. Application of 10 nM PACAP-27 to the cell from a puffer pipette caused a decrease in membrane conductance, similar to the effects of the related peptide, GLP-1, which was ascribed to the closure of ATP-sensitive  $K^+$ -channels (8).

Interestingly, PACAP produced no rise in  $[Ca^{2+}]_i$  in the presence of  $Ca^{2+}_o$ , but in the absence of  $Na^+_o$  (Fig. 4B). A similar dependence on  $Na^+_o$  has also been demonstrated for the PACAP-related peptide, GLP-1, in rat islets (18). The  $Na^+_o$  dependence of the rise in  $[Ca^{2+}]_i$  in response to PACAP is likely to be explained at least in part by the inhibition of the inward current, which has a large  $Na^+$  component (see below, Fig. 6A).

$La^{3+}$  is frequently used as a blocker of a wide range of voltage-dependent and -independent  $Ca^{2+}$  channels. If the effect of PACAP on  $[Ca^{2+}]_i$  results from activation of such channels, then  $La^{3+}$  might be expected to inhibit the PACAP-induced inward current. To test this possibility, HIT-T15 cells were held under voltage clamp at  $-70$  mV, and  $\pm 10$  mV voltage steps were applied to monitor membrane conductance. An inward current, increase in membrane conductance (Fig. 5A), and rise in  $[Ca^{2+}]_i$  (Fig. 5B) were observed in response to PACAP-27. Direct application of 100  $\mu$ M  $La^{3+}$  from a puffer pipette caused a reversible decrease in both the inward current and membrane conductance (Fig. 5A), indicating that at least one component of the current is blocked by  $La^{3+}$ . The PACAP-27-activated current was not completely blocked by  $La^{3+}$ , which did, however, completely block the rise in  $[Ca^{2+}]_i$  in response to a voltage step from  $-70$  to 0 mV (data not shown), indicating that VDCCs were fully blocked.

The current activated by PACAP is also activated by forskolin, 8Br-cAMP, and Sp-cAMPs. To obtain an estimate for the reversal potential of the current, cells were dialyzed with  $Cs^+$  (substituted for  $K^+$  in the pipette solution) in the perforated patch configuration. Voltage ramps of 100-msec duration from  $-70$  to 30 mV were applied, and the currents in response to these ramps before stimulation with 8Br-cAMP were subtracted from currents during the response to stimulation with 8Br-cAMP to obtain the current-voltage relation of the evoked current. These currents had an apparent reversal potential of  $-28.0 \pm 2.3$  mV ( $n = 4$ ). This value for the

FIG. 3. Simultaneous records of  $[Ca]_i$  and membrane potential or membrane current. Records from a single HIT cell bathed in 2 mM glucose. Initially, the fura-2-loaded cell was under perforated patch current clamp and responded to the application of 10 nM PACAP-27 (P27) by depolarizing and firing action potentials (A), accompanied by a rise in  $[Ca^{2+}]_i$  (B). When the same cell was voltage clamped at  $-70$  mV, application of PACAP-27 induced a large inward current (C) and a rise in  $[Ca^{2+}]_i$  (D).



reversal potential is consistent with the activation of non-specific cation channels.

In the presence of physiological ion gradients, the main charge carrier through such nonspecific cation channels would be expected to be  $Na^+$ . A HIT-T15 cell stimulated with 1 mM 8Br-cAMP developed a large inward current and an increase in membrane conductance (Fig. 6A). During the response, an extracellular solution in which  $Na^+$  had been substituted by NMG was applied, which substantially and reversibly reduced the current and membrane conductance (Fig. 6A). The current evoked by 8Br-cAMP could also be inhibited by  $La^{3+}$  (Fig. 6B).

PACAP produced a marked increase in the  $Mn^{2+}$  permeability of HIT cells (Fig. 7). A 60-sec pulse of  $Mn^{2+}$  ( $Mn^{2+}$  was substituted for  $Ca^{2+}$  of normal bath solution) applied to a single cell from a puffer pipette had no effect on the 350/380 nm fluorescence ratio (Fig. 7A) or raw fluorescence intensity (Fig. 7B) before stimulation of the cell with PACAP-38. Stimulation with PACAP induced a rise in  $[Ca^{2+}]_i$  (increased fluorescence ratio), and the subsequent application of a 60-sec pulse of  $Mn^{2+}$  led to quenching of the fura-2 fluorescence (Fig. 7B).  $Mn^{2+}$ -permeant channels believed to be associated with capacitative  $Ca^{2+}$  entry ( $I_{CRAC}$ ) have been reported in HIT-T15 cells (14), RINm5F cells (19), and mouse  $\beta$ -cells (20). However, it remains to be determined whether the activation of an  $I_{CRAC}$  explains the increased  $Mn^{2+}$  permeability observed after treatment of HIT cells with PACAP.

## Discussion

In this report we demonstrate that in HIT-T15 cells, the biologically active isoforms of PACAP induce membrane depolarization through at least two distinct mechanisms. First, PACAP reduces the membrane conductance and increases membrane excitability in a manner similar to that previously described for GLP-1 (11). This effect most likely results from a potentiation by PACAP of the inhibitory effect of glucose on ATP-K channels. Secondly, PACAP increases membrane conductance and activates a pronounced inward current that is predominantly a  $Na^+$  current, blockable by  $La^{3+}$ , and exhibits a reversal potential of about  $-28$  mV. We propose that the inward current results from activation of ion channels similar to the cAMP- and  $Ca^{2+}$ -gated nonspecific cation channels (Ca-NS channels) reported in CRI-G1 insulinoma cells (9).

PACAP also induced a rise in  $[Ca^{2+}]_i$  by several distinct mechanisms. One process results from the effects on ATP-K and Ca-NS channels, as described above, leading to membrane depolarization and activation of L-type VDCCs. The  $Ca^{2+}$  current through VDCCs may also be enhanced by cAMP-dependent protein phosphorylation (12). A second mechanism by which PACAP raised  $[Ca^{2+}]_i$  was observed under conditions in which HIT-T15 cells were voltage clamped at  $-70$  mV (and also  $-120$  mV; data not shown), thereby preventing the opening of VDCCs (17). Although the threshold for activation of VDCCs can shift to more negative membrane potentials under conditions of pharmacological




FIG. 4. The PACAP-induced rise in  $[Ca^{2+}]_i$  is dependent on extracellular  $Ca^{2+}$  and  $Na^+$ . A, The rise in  $[Ca^{2+}]_i$  of a HIT cell in response to a 10-sec pulse of 2 nM PACAP-38 is reversed by switching the bath solution to one that is nominally calcium free. Returning the cell to a bath solution containing  $Ca^{2+}$  caused  $[Ca^{2+}]_i$  to rise again, indicating that the conductance mechanism remains open. All solutions contained 0.8 mM glucose. Similar results were obtained in all five cells tested. B, The sodium dependence of PACAP. A HIT cell was bathed in sodium free (NMG-substituted) saline and initially showed no rise in  $[Ca^{2+}]_i$  in response to a 10-sec pulse of 2 nM PACAP 38 (P38) in Na-free NMG saline. In the continued presence of Na-free bathing saline, the same cell showed a large rise in  $[Ca^{2+}]_i$  in response to a 10-sec pulse of 2 nM PACAP 38 in sodium-containing (138 mM  $Na^+$ ) saline. All solutions contained 0.8 mM glucose. Similar effects were seen in all five cells tested using the same protocol.

stimulation (21), these shifts would not be expected to be large enough to significantly activate VDCCs at  $-70$  mV. Therefore, the ability of PACAP to raise  $[Ca^{2+}]_i$  under conditions of voltage clamp indicates the existence of a  $Ca^{2+}$  signaling mechanism involving voltage-independent  $Ca^{2+}$  channels and/or the release of  $Ca^{2+}$  from intracellular stores. These observations contrast with a previous report suggesting that the mechanism by which PACAP raises  $[Ca^{2+}]_i$  in  $\beta$ -cells is by increasing the activity of L-type VDCCs (4).

We observed that PACAP was without effect on  $[Ca^{2+}]_i$  under conditions in which the extracellular solution contained  $Ca^{2+}$ , but not  $Na^+$ . This suggests that mobilization of intracellular  $Ca^{2+}$  by PACAP might be initiated by the entry of  $Na^+$  through Ca-NS channels and/or effects of cAMP on organellar  $Ca^{2+}$  stores. The elevation of  $[Na^+]_i$  in  $\beta$ -cells is known to trigger  $Ca^{2+}$  release from intracellular stores (22).




FIG. 5. PACAP-27 increases membrane conductance, which is blocked by  $La^{3+}$ . A HIT cell was held under perforated patch voltage clamp at  $-70$  mV, and  $\pm 10$  mV steps were applied to monitor membrane conductance. Application of 10 nM PACAP-27 to the cell induced an inward current and increased the membrane conductance, both of which were reduced by  $La^{3+}$  (A). The inward current was accompanied by a rise in  $[Ca^{2+}]_i$  (B).  $[Ca^{2+}]_i$  started to recover toward normal levels after the application of  $La^{3+}$  despite incomplete block of the PACAP-induced current and conductance increase.

It has also been demonstrated that the secretory granules form a storage site for intracellular  $Ca^{2+}$ , and that this  $Ca^{2+}$  pool is rapidly altered by stimulation with glucose and by intracellular factors, notably cAMP and  $Na^+_i$  (23). Furthermore, cAMP reportedly stimulates  $^{45}Ca^{2+}$  efflux from  $\beta$ -cell secretory granules (24). Therefore, ample precedent exists supporting the concept of an important influence of  $Na^+$  and cAMP on  $\beta$ -cell  $Ca^{2+}$  signaling systems unrelated to VDCCs. Although the mechanism by which cAMP mobilizes  $Ca^{2+}$  in  $\beta$ -cells remains to be determined, cAMP-dependent phosphorylation of the inositol trisphosphate ( $IP_3$ ) receptor by PKA can increase  $Ca^{2+}$  release (25, 26). However, there is at present no direct evidence that the type 3  $IP_3$  receptor, located in the membrane of insulin secretory granules (27), is a substrate for PKA (28).

We also report that stimulation of HIT-T15 cells by PACAP results in the activation of a  $Mn^{2+}$ -permeant divalent cation influx pathway. This was demonstrated by the quenching of intracellular fura-2 fluorescence by externally applied  $Mn^{2+}$ . Quenching is unlikely to reflect entry of  $Mn^{2+}$  via VDCCs, as it is well established that  $Mn^{2+}$  is a broad spectrum blocker of this class of  $Ca^{2+}$  channels. Instead, we propose that the entry of  $Mn^{2+}$  reflects the activation of a  $Ca^{2+}$ -release-activated  $Ca^{2+}$ -current ( $I_{CRAC}$ ) (14, 19). This is consistent with the proposed ability of PACAP to mobilize intracellular  $Ca^{2+}$




FIG. 6. 8Br-cAMP activates a current permeant to Na<sup>+</sup> and blocked by La<sup>3+</sup>. A HIT cell bathed in 0.8 mM glucose saline was held at -70 mV in perforated patch voltage clamp, with 750-msec steps of ±10 mV applied. The cell initially had a very low membrane conductance until it was stimulated with 1 mM 8Br-cAMP (indicated by bar). A large inward current and increase in membrane conductance were observed in response to 8Br-cAMP (A). During the response, a pulse of Na<sup>+</sup>-free saline (substituted by NMG) was applied (bar), and current and conductance were both reduced (A), suggesting that a component of the current was carried by Na<sup>+</sup>. A different HIT cell, under the same recording conditions as those defined in A, was stimulated by 8Br-cAMP, generating an inward current and increase in conductance, which was inhibited by the application of 250 μM La<sup>3+</sup> (B).




FIG. 7. Activation of a Mn<sup>2+</sup>-permeant conductance after PACAP stimulation. A HIT cell, bathed in 0.8 mM glucose saline, showed no quenching of fura-2 fluorescence during a 60-sec pulse of Mn<sup>2+</sup> before stimulation with a 10-sec pulse of 2 nM PACAP-38 (P38), but showed marked quenching after PACAP stimulation. Periods of application are indicated by bars. The Mn<sup>2+</sup>-saline was applied from a puffer pipette and prepared by substitution of Mn<sup>2+</sup> for Ca<sup>2+</sup> (2.6 mM).

stores, because the depletion of such stores is known in some cell types to activate I<sub>CRAC</sub>.

The PACAP-R3, present in HIT cells, couples to phospholipase-C, generation of IP<sub>3</sub>, and activation of protein kinase-C (PKC) when expressed in oocytes (5). Although such coupling has yet to be demonstrated for the native receptor in β-cells or insulinoma cells, it remains possible that an effect of PACAP on IP<sub>3</sub> production or PKC activity contributes to the observed rise in [Ca<sup>2+</sup>]<sub>i</sub> reported here. Activation of I<sub>CRAC</sub> in RINm5F cells has been observed after stimulation of PKC (19), and PKC can also phosphorylate IP<sub>3</sub> receptors (29). Hence, regulation of intracellular Ca<sup>2+</sup> stores and activation of I<sub>CRAC</sub> may represent an important point of convergence, at which signal transduction cross-talk occurs between the cAMP and PLC/PKC/IP<sub>3</sub> signaling systems in β-cells.

The physiological role of PACAP remains to be fully clarified. PACAP immunoreactivity has been observed in nerve endings within the pancreas and was occasionally found within the islets (3). These observations suggest that PACAP may have local paracrine effects *in vivo*. We propose that a neuro-entero-endocrine loop may exist whereby feeding stimulates the local release of PACAP within pancreatic islets, thereby triggering insulin secretion. This release would

provide a rapid augmentation of insulin secretion before the intestinal incretins come into play.

In summary, both isopeptides of PACAP cause the voltage-independent activation of an inward current, which we suggest is carried through Ca-NS channels. PACAP also decreases the resting membrane conductance, probably through inhibition of ATP-K channels. These effects result in depolarization of the cell and activation of L-type VDCCs. Furthermore, there are additional mechanisms by which  $[Ca^{2+}]_i$  rises in a membrane potential-independent manner.  $Ca^{2+}$  entry through voltage-independent  $Ca^{2+}$  channels, including  $I_{CRAC}$ , may contribute to the rise in  $[Ca^{2+}]_i$ , and the elevation of intracellular cAMP and/or  $Na^+$  levels may stimulate  $Ca^{2+}$  release from intracellular stores. Acting in concert, these various  $Ca^{2+}$  signaling pathways are expected to exert important stimulatory influences on pancreatic insulin secretion.

### References

- Miyata A, Arimura A, Dahl RR, Minamino N, Vehara A, Jiang L, Culler MD, Coy DH 1989 Isolation of a novel 38 residue-hypothalamic polypeptide which stimulates adenylate cyclase in pituitary cells. *Biochem Biophys Res Commun* 164:567-574
- Kimura C, Ohkubo S, Og K, Hosoya M, Itoh Y, Onda H, Miyata A, Jiang L, Dahl RR, Stibbs HH, Arimura A, Fujino M 1990 A novel peptide which stimulates adenylate cyclase: molecular cloning and characterization of the ovine and human cDNAs. *Biochem Biophys Res Commun* 166:81-89
- Fridolf T, Sundler F, Ahren B 1992 Pituitary adenylate cyclase-activating polypeptide (PACAP): occurrence in rodent pancreas and effects on insulin and glucagon secretion. *Cell Tissue Res* 269:275-279
- Yada T, Sakurada M, Ihida K, Nakata M, Murata F, Arimura A, Kikuchi M 1994 Pituitary adenylate cyclase-activating polypeptide is an extraordinarily potent intra-pancreatic regulator of insulin secretion from islet  $\beta$ -cells. *J Biol Chem* 269:1290-1293
- Inagaki N, Yoshida H, Mizuta M, Mizuno N, Fujii Y, Gono T, Miyazaki J-I, Seino S 1994 Cloning and functional characterization of a third pituitary adenylate cyclase-activating polypeptide receptor subtype expressed in insulin-secreting cells. *Proc Natl Acad Sci USA* 91:2679-2683
- Lutz EM, Sheward WJ, West KM, Morrow JA, Fink G, Harmar AJ 1993 The  $VIP_2$  receptor: molecular characterisation of a cDNA encoding a novel receptor for vasoactive intestinal peptide. *FEBS Lett* 334:3-8
- Schomerus E, Poch A, Bunting R, Mason WT, McArdle CA 1994 Effects of pituitary adenylate cyclase-activating polypeptide in the pituitary: activation of two signal transduction pathways in the gonadotrope-derived  $\alpha T3-1$  cell line. *Endocrinology* 134:315-323
- Yokota C, Kawai K, Ohashi S, Susuki S, Yamashita K 1993 Stimulatory effects of pituitary adenylate cyclase-activating polypeptide (PACAP) on insulin and glucagon release from the isolated perfused rat pancreas. *Acta Endocrinol (Copenh)* 129:473-479
- Reale V, Hales CN, Ashford MLJ 1994 Nucleotide regulation of a calcium-activated cation channel in the rat insulinoma cell line, CRI-G1. *J Membr Biol* 141:101-112
- Sturgess NC, Hales CN, Ashford MLJ 1986 Inhibition of a calcium-activated non-selective cation channel, in a rat insulinoma cell line, by adenine derivatives. *FEBS Lett* 208:397-400
- Holz GG, Kuhlreier WM, Habener JF 1993 Pancreatic  $\beta$ -cells are rendered glucose competent by the insulinotropic hormone GLP-1(7-37). *Nature* 361:362-365
- Ammala C, Ashcroft FM, Rorsman P 1993 Calcium-independent potentiation of insulin release by cyclic AMP in single  $\beta$ -cells. *Nature* 363:356-358
- Leech CA, Holz GG 1994 Application of patch clamp methods to the study of calcium currents and calcium channels. *Methods Cell Biol* 40:135-151
- Leech CA, Holz GG, Habener JF 1994 Voltage-independent calcium channels mediate slow oscillations of cytosolic calcium that are glucose dependent in pancreatic  $\beta$ -cells. *Endocrinology* 135:365-372
- Grynkiewicz G, Poenie M, Tsien RY 1985 A new generation of  $Ca^{2+}$  indicators with greatly improved fluorescence properties. *J Biol Chem* 260:3440-3450
- Horn R, Marty A 1988 Muscarinic activation of ionic currents by a new whole-cell recording method. *J Gen Physiol* 92:145-159
- Hiriart M, Matteson DR 1988  $Na^+$  channels and two types of  $Ca^{2+}$  channels in rat pancreatic  $\beta$ -cells identified with the reverse hemolytic plaque assay. *J Gen Physiol* 91:617-639
- Fridolf T, Ahren B 1993 Effects of glucagon-like peptide-1(7-36)amide on the cytoplasmic  $Ca^{2+}$ -concentration in rat islet cells. *Mol Cell Endocrinol* 96:85-90
- Bode H-P, Goke B 1994 Protein kinase C activates capacitance calcium entry in the insulin secreting cell line RINm5F. *FEBS Lett* 339:307-311
- Worley JF, McIntyre MS, Spencer B, Mertz RJ, Roe MW, Dukes ID 1994 Endoplasmic reticulum calcium store regulates membrane potential in mouse islet  $\beta$ -cells. *J Biol Chem* 269:14359-14362
- Velasco JM, Petersen JUH, Petersen OH 1988 Single-channel  $Ba^{2+}$  currents in insulin-secreting cells are activated by glyceraldehyde stimulation. *FEBS Lett* 123:366-370
- Herchuelz A, Malaisse WJ 1980 Regulation of calcium fluxes in rat pancreatic islets: dissimilar effects of glucose and of sodium ion accumulation. *J Physiol* 302:263-280
- Wolters GHJ, Pasma A, Wiegman JB, Konijnendijk W 1984 Glucose-induced changes in histochemically determined  $Ca^{2+}$  in  $\beta$ -cell granules,  $^{45}Ca$  uptake, and total  $Ca^{2+}$  of rat pancreatic islets. *Diabetes* 33:409-414
- Hahn HJ, Gylfe E, Hellman B 1980 Calcium and pancreatic  $\beta$ -cell function. VII. Evidence for cyclic AMP-induced translocation of intracellular calcium. *Biochim Biophys Acta* 630:425-432
- Burgess GM, Bird GSTJ, Obie JF, Putney JW 1991 The mechanism for synergism between phospholipase C- and adenylyl cyclase-linked hormones in the liver. Cyclic AMP-dependent kinase augments inositol trisphosphate-mediated  $Ca^{2+}$  mobilization without increasing the cellular levels of inositol polyphosphates. *J Biol Chem* 266:4772-4781
- Nakade S, Rhee SK, Hamanaka H, Mikoshiba K 1994 Cyclic AMP-dependent phosphorylation of an immunopurified homotetrameric inositol 1,4,5-trisphosphate receptor (type I) increase  $Ca^{2+}$  flux in reconstituted lipid vesicles. *J Biol Chem* 269:6735-6742
- Blondel O, Moody MM, Depaoli AM, Sharp AH, Ross CA, Swift H, Bell GI 1994 Localization of inositol trisphosphate receptor subtype 3 to insulin and somatostatin secretory granules and regulation of expression in islets and insulinoma cells. *Proc Natl Acad Sci USA* 91:7777-7781
- Blondel O, Takeda J, Janssen H, Seino S, Bell GI 1994 Sequence and functional characterization of a third inositol trisphosphate receptor subtype, IP3R-3, expressed in pancreatic islets, kidney, gastrointestinal tract, and other tissues. *J Biol Chem* 268:11356-11363
- Ferris CD, Haganir RL, Bredt DS, Cameron AM, Snyder SH 1991 Inositol trisphosphate receptor: phosphorylation by protein kinase C and calcium calmodulin-dependent protein kinases in reconstituted lipid vesicles. *Proc Natl Acad Sci USA* 88:2232-2235