

Characterization of the Electrically Evoked Release of Substance P from Dorsal Root Ganglion Neurons: Methods and Dihydropyridine Sensitivity

George G. Holz IV, Kathleen Dunlap, and Richard M. Kream¹

Departments of Physiology and ¹Anesthesiology, Tufts University School of Medicine and New England Medical Center, Boston, Massachusetts 02111

The mechanism by which dihydropyridines (DHPs) modulate the electrically evoked or KCl-induced release of substance P (SP) from embryonic chick dorsal root ganglion (DRG) neurons was investigated in the present study. The release of SP, as measured by radioimmunoassay (RIA), was characterized in terms of its dependence on extracellular calcium ion, its stimulus-response relationship, its sensitivity to the calcium-channel blocker omega conus toxin (ω -CgTx), and its modulation by the DHPs Bay K 8644 and nifedipine. Here it is reported that ω -CgTx (1 μ M) blocked the electrically evoked release of SP. In contrast, the calcium-channel agonist Bay K 8644 (5 μ M) facilitated the release of SP (by 45%), whereas the calcium-channel antagonist nifedipine (5 μ M) was without effect. When the release of SP was triggered by depolarization of cultures with 60 mM KCl, the actions of the DHPs became much more pronounced. Under these conditions, Bay K 8644 facilitated (by 115%), whereas nifedipine inhibited (by 58%), peptide secretion. Voltage-clamp analysis of DRG cell calcium currents demonstrated that these actions of ω -CgTx, Bay K 8644, and nifedipine are explicable in terms of their effects on the slowly inactivating (L-type) calcium current. On the basis of these findings, it is suggested that the SP release mechanism exhibits DHP sensitivity due to the involvement of L-type calcium channels in the neurosecretory process. This model predicts that the voltage and time-dependent antagonist actions of nifedipine are sufficient to explain its failure to inhibit the electrically evoked release of SP.

Substance P (SP) is a tachykinin neuropeptide synthesized and secreted by finely myelinated and unmyelinated primary afferent neurons (Pernow, 1983a). Subsequent to its isolation, purification, and chemical characterization by Chang and Leeman in 1970, attention focused on the role of SP as an excitatory transmitter or modulator at spinal terminations of primary afferent nociceptors (Otsuka and Konishi, 1976; Henry, 1977, 1982; Jessel and Iversen, 1977; Gamse et al., 1979; Nicoll et al., 1980; Yaksh et al., 1980). More recently, attention has also been fo-

cused on the peripheral terminations of primary afferent neurons, where released SP is implicated as a mediator of neurogenic inflammation (Pernow, 1983b, 1985), and as a modulator of immune system function (Goetzl et al., 1985; Payan and Goetzl, 1985). Despite the apparent importance of these diverse biological actions of SP, very little is known regarding cellular processes that regulate its release.

In an attempt to address this question, we have studied the mechanism by which depolarization induces the release of SP from embryonic chick dorsal root ganglion (DRG) neurons. It was previously reported that these primary afferent neurons synthesize and secrete SP when grown in dissociated cell culture (Mudge, 1979, 1981; Mudge et al., 1979). These earlier studies examined the mechanism by which depolarizing concentrations of KCl induce peptide secretion. In the present paper we characterize the electrically evoked release of SP, as measured by RIA and as authenticated by high-pressure liquid chromatography (HPLC). In addition, we have begun to characterize the type of calcium channel(s) through which calcium enters to trigger secretion.

Recent electrophysiological studies have determined that 3 types of depolarization-induced calcium current may be recorded from the soma membrane of DRG neurons (Nowycky et al., 1985a). These 3 currents presumably reflect calcium influx through 3 distinct subtypes of voltage-dependent calcium channel. Current flow through one of these, termed the L-type channel, is selectively inhibited by the dihydropyridine (DHP) calcium-channel antagonist nifedipine (Rane et al., 1987), and facilitated by the DHP calcium-channel agonist Bay K 8644 (Nowycky et al., 1985a, b). Moreover, Perney et al. (1986) reported that antagonist DHPs block the KCl-induced release of SP from neonate rat DRG cell cultures. In order to more fully investigate the potential role of L-type calcium channels in neurosecretory processes, we have examined whether DHPs also modulate the electrically evoked release of SP.

We now report that Bay K 8644 facilitated both the electrically evoked and KCl-induced release of SP. In contrast, nifedipine failed to inhibit release evoked by electrical-field stimulation, but reduced peptide secretion induced by 60 mM KCl. These findings are interpreted in terms of the voltage and time-dependent actions of DHPs on L-type calcium channels, as assessed by electrophysiological recordings from these same cells. A model is proposed whereby neuropeptide secretion is triggered by calcium influx through L-type calcium channels located at release sites on primary afferent nerve terminals. This model predicts that the voltage and time-dependent antagonist actions

Received Dec. 9, 1986; revised May 26, 1987; accepted July 28, 1987.

This work was supported by grants to K.D. from the NIH (NS 16438), the Klingenstein Fund, and the American Heart Association (with funds contributed, in part, by the Massachusetts affiliate), and to R.M.K. from NIDA (DA 04128) and the New England Medical Center (BRSG 19-807419).

Correspondence should be addressed to Dr. George G. Holz IV, Department of Physiology, Tufts University School of Medicine, 136 Harrison Avenue, Boston, MA 02111.

Copyright © 1988 Society for Neuroscience 0270-6474/88/020463-09\$02.00/0




Figure 1. Electrical field stimulation of DRG cell cultures. *A*, Design of the stimulation chamber. Bipolar platinum steel electrodes are inserted into the 60 mm tissue culture dish so that they contact the solution bathing the cells. Note that when inserted, each electrode occupies one-third of the inner circumference of the dish. *B*, *C*, Action potentials (note different time scales) recorded from the same DRG neuron in response to electrical field stimulation (square-wave DC pulses, 110 V, 3 msec duration). A single action potential is illustrated in *B* (arrowhead indicates stimulus artifact), whereas a train of action potentials evoked by stimulation at 25 Hz is illustrated in *C*. The resting potential was -58 mV. The horizontal line through the spikes indicates the zero potential determined after withdrawing the electrode from the cell. The recording solution consisted of HBS containing 2 mM Ca^{2+} and 1 mM Ba^{2+} . Calibration bar: 50 mV, 10 msec (*B*); 50 mV, 40 msec (*C*).

of nifedipine are sufficient to explain its failure to inhibit the electrically evoked release of SP.

Materials and Methods

Preparation of DRG cell cultures. Primary cultures of embryonic chick DRG cells were prepared as previously described (Dichter and Fischbach, 1977; Mudge et al., 1979; Dunlap and Fischbach, 1981). DRGs from 11- or 12-d-old embryos were incubated for 45 min in Ca^{2+} - and Mg^{2+} -free Puck's solution, resuspended in Eagle's Minimal Essential Medium (MEM, supplemented with 7 S nerve growth factor, 5% chick embryo extract, 10% horse serum, 1 mM glutamine, 50 U/ml penicillin, and 50 $\mu\text{g}/\text{ml}$ streptomycin), and mechanically dissociated by trituration to give a single-cell suspension. The DRG cells were gamma-irradiated (5000 rads) and plated on 60 mm collagen-coated tissue culture dishes at a density of 150,000–200,000 neurons/dish. This density is 2–2.5 times that used in previous studies examining the electrophysiological properties of DRG cells (Dunlap and Fischbach, 1978, 1981). Culture medium was changed every 2–3 d beginning on day 3. The peak phase of non-neuronal cell death was observed on days 3–5.

The plating density was a critical factor in these experiments. Lower plating densities resulted in poor cell adhesion and limited neurite outgrowth. Large numbers of non-neuronal cells were observed with higher plating densities. In both cases the cellular content of SP was reduced to less than 1 ng/culture, whereas a cellular content of 10–15 ng/culture was observed by day 16 in those cultures that we judged to be of optimal density. Highest levels of SP were observed in cultures exhibiting dense patterns of neurites, clear cell bodies with prominent nucleoli, and few non-neuronal cells. In those cultures the cells remained firmly attached

to the collagen substrate even with repeated exchanges of the bathing solution.

Stimulation protocol for substance P release experiments. DRG cell cultures were stimulated under sterile conditions at room temperature using bipolar platinum steel electrodes. This allowed repeated use of the same cultures for experiments on as many as 10 different days. As illustrated in Figure 1*A*, 2 electrodes were positioned opposite each other on the outer circumference of a plastic insert that fit snugly against the inner circumference of the culture dish. Each electrode occupied one-third of the circumference of the dish, and a roof over the stimulation chamber protected against drying of the chamber contents. Stimuli (square-wave DC pulses, 3 msec duration, 110 V) were delivered at 1 Hz, and the stimulus polarity was reversed every 30 sec. Microelectrode recordings demonstrated that field stimulation generated action potentials of normal configuration in all cells tested (Fig. 1, *B*, *C*).

Composition of solutions. All solutions used in the release experiments consisted of saline buffered with 25 mM HEPES, pH 7.4. The HEPES-buffered saline (HBS) contained (in mM): 132 NaCl, 2.5 KCl, 2.0 CaCl_2 , 0.8 MgCl_2 , and 0.04% BSA. For solutions containing 1 mM Ba^{2+} or 5 mM Co^{2+} , an appropriate volume of concentrated BaCl_2 or CoCl_2 stock solution was added directly to the HBS. Nifedipine was obtained from Sigma, Bay K 8644 from Miles Laboratories, and ω -CgTx from Peninsula Laboratories. Nifedipine and Bay K 8644 were prepared as concentrated (5 mM) stock solutions in absolute ethanol. Stock solutions were diluted in HBS to give 5 μM test solutions (final concentration of ethanol, 0.10%). In all experiments examining modulation of release by dihydropyridines, control solutions (i.e., no drugs added) contained 0.10% ethanol.

Radioimmunoassay for released substance P. Field stimulation-in-

duced release of SP was measured by direct radioimmunoassay (RIA) of the solution bathing the cells. The RIA employed an antibody specific for authentic SP (1-11) and its sulfoxide derivative, with less than 0.01% cross-reactivity to the related tachykinins physalaemin and substance K (Kream et al., 1985). Tracer was synthesized by radioiodination of the Bolton-Hunter conjugate of SP. The specific activity of reverse-phase HPLC-purified tracer was approximately 2000 Ci/mmol.

Prior to electrical field stimulation, each culture was bathed in 1.7 ml of the solution to be tested. Following field stimulation, a 1.64 ml aliquot of the bathing solution was removed for RIA. To each 1.64 ml test sample was added 60 μ l of RIA buffer, containing (in gm/liter): 44 NaCl, 10 BSA, 20 peptone, 0.2 cytochrome C, 3.7 disodium EDTA, 0.1 sodium azide, 60 HEPES, 0.01 aprotinin, 0.25 leupeptin, 25 pepstatin-A, and 0.1 phenylmethylsulfonyl fluoride, pH 8.0. Aliquots (0.4 ml) of this HBS/RIA buffer solution were assayed in quadruplicate. To each aliquot was added 20 μ l of SP antibody such that the final antibody dilution factor was 150,000–250,000-fold (33–45% of the immunoreactive tracer was bound in this range of dilutions). Next, 20 μ l of 125 I-SP tracer was added 16 hr following addition of the antibody. The assay was equilibrated for 24 hr, and bound tracer was separated from free tracer by precipitation and centrifugation of the antibody-tracer complex in 2 ml 100% ethanol.

The sensitivity of the RIA was adjusted by varying the concentration of SP antibody. In most experiments, the IC_{50} value for displacement of immunoreactive tracer by nonradioactive SP was 125 pg SP/assay and the lower limit of detection was 16 pg SP/assay. Sensitivity as low as 2 pg SP/assay was obtained with more dilute antibody, but this level of detection was not required in most experiments. Standard curves were generated by assaying serial dilutions of synthetic SP standard (Sigma) that was diluted in the same HBS test solution as was used for the release experiment. Standard curves were linearized by logit-log transformation of these data. None of the drugs tested interfered with binding of the tracer to the antibody. The intra-assay coefficient of variation was less than 10%.

Authentication of substance P by HPLC. Authentication of SP extracted from DRG cell cultures was performed by reverse-phase HPLC as previously described (Kream et al., 1985). SP was extracted in 2 ml of 6 N guanidine HCl and the extract subjected to an initial purification (step 1) by reverse-phase chromatography on Baker 10 SPE octadecyl C18 columns. SP was eluted in 2 ml of 50% CH_3CN containing 0.1% CF_3COOH . Aliquots (50 μ l) were lyophilized, resuspended in RIA buffer, and assayed by RIA to determine the total cellular content of SP per culture.

HPLC was performed using a 3.9×150 mm Waters C18 Nova-Pak column. The eluate from step 1 was diluted 10-fold with distilled water, and 1 ml aliquots of this solution were serially loaded onto the column. Elution was initially achieved under isocratic conditions (8 min), with a mobile phase of 15% (wt/vol) CH_3CN in 0.1% CF_3COOH , followed by a linear 15–45% gradient of CH_3CN in 0.1% CF_3COOH (30 min). The flow rate was 1 ml/min, and 0.5 ml fractions were collected. Aliquots (50 μ l) of each fraction were lyophilized, resuspended in buffer, and assayed for SP by RIA. Recovery of SP typically was 85–90%.

Intracellular recordings from DRG cell bodies. Intracellular microelectrode and voltage-clamp recordings of DRG cell action potentials and calcium currents were obtained as previously described (Dichter and Fischbach, 1977; Rane and Dunlap, 1986). In both recording configurations, the extracellular solution was the same as that used in the release experiments (the divalent cation concentrations were 2 mM Ca^{2+} and 1 mM Ba^{2+}), except that 0.3 μ M tetrodotoxin was added to the solution when recording calcium currents. For whole-cell patch-clamp recordings of calcium currents, the pipette solution contained (in mM): 150 CsCl, 5 bis(*o*-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid (BAPTA), 5 MgATP, and 10 HEPES (pH 7.3). All drugs were applied by pressure-ejection from a blunt-tipped micropipette.

Results

Authentication of substance P immunoreactivity

Release experiments were performed using 9–16-d-old DRG cell cultures. In preliminary experiments it was demonstrated that electrical field stimulation at 1 Hz for 30 sec resulted in the appearance of detectable levels of SP immunoreactivity in the bathing solution. SP synthesized by DRG cells was judged to be authentic SP on the basis of chromatographic and im-




Figure 2. Authentication of SP immunoreactivity by HPLC. The cellular content of SP from 6 cultures was extracted in 6 N guanidine HCl and fractionated by HPLC as described in Materials and Methods. Fractions (0.5 ml) were collected and 20 μ l aliquots assayed for SP by RIA. The elution profile of DRG cell SP is indicated by the cross-hatched columns. The arrowheads indicate the position determined for the major immunoreactive fractions of synthetic SP and the sulfoxide derivative of SP (SP-O).

munoreactive properties. DRG cell extracts were fractionated by reverse-phase HPLC, and the elution profile of cellular SP was monitored by RIA. As illustrated in Figure 2, the retention time for cellular SP matched that of synthetic SP standard. In addition, a smaller peak corresponding to the sulfoxide derivative (SP-O, to which the antibody cross-reacts; Kream et al., 1985) of SP was also noted.


As expected, SP extracted, or released, from DRG cell cultures displayed a dose-dependent displacement of the binding of 125 I-SP to the SP antibody. As illustrated in Figure 3, a standard curve was generated by assaying serial dilutions of synthetic SP. When serial dilutions of cellular extract (Fig. 3, square symbols) or released SP (circles) were assayed, the displacement curves generated displayed slopes identical to the standard curve (triangles). For these reasons, the immunoreactivity recognized by our antibody will henceforth be referred to as SP.

Characterization of the effects of divalent cations on field stimulation-induced release of substance P

We characterized the effects of electrical field stimulation on the SP release process. As illustrated in Figure 4, when cultures were bathed in HBS containing 3 mM Ca^{2+} , stimulation (5 min at 1 Hz) increased SP levels in the bathing solution 4.5-fold over baseline values (unstimulated: 68 ± 16 pg/culture; stimulated: 305 ± 26 pg/culture; mean \pm SEM; $n = 3$). The amount of SP released was 2.3% of the total cellular content of SP per culture (total cellular content was calculated as the sum of the amount released plus the amount remaining in the cells).

The release of SP was calcium-dependent: it was blocked by HBS containing 0.1 mM Ca^{2+} and 5.0 mM Co^{2+} (Fig. 4). In contrast, a marked facilitation of release was observed when the HBS contained 1 mM Ba^{2+} and 2 mM Ca^{2+} . In fact, Ba^{2+} more than doubled the amount of SP released (Fig. 4). Note that in the presence of Ca^{2+} , 1 mM Ba^{2+} was not a secretagogue: baseline levels of SP were not elevated relative to control. Intracellular

Figure 3. Parallel displacement curves generated for synthetic, cellular, and released SP. Displacement curves were linearized by logit-log transformation of the binding data. A standard curve was generated by assaying serial dilutions of synthetic SP (triangles, upper abscissa). Cellular and released SP were subjected to an initial purification on Baker 10 SPE C18 columns, as described in Materials and Methods. SP was eluted in 2 ml 50% acetonitrile. Serial dilutions of the eluates were assayed for SP by RIA, and displacement curves generated for cellular (squares) and released (circles) SP. The lower abscissa indicates the volume (in μ l) of the original 2 ml eluate that was assayed by RIA.



recordings demonstrated that the action of Ba^{2+} can be explained by its effect on the duration of DRG cell action potentials. As illustrated in the inset of Figure 4, these action potentials exhibit a plateau phase that is reduced in duration by Co^{2+} and increased in duration by Ba^{2+} (Dichter and Fischbach, 1977). Since the duration of the plateau phase is directly related to the amount of Ca^{2+} that enters with each action potential, it is likely that Ba^{2+} facilitates the release of SP by increasing spike duration.

Determination of the stimulus-response relationship


Figure 5 illustrates the stimulus-response relationship for field stimulation-induced release of SP over a range of 60–600 stimuli delivered at a frequency of 1 Hz. In these experiments the HBS contained 1 mM Ba^{2+} and 2 mM Ca^{2+} in order to maximize the amount of SP released. Baseline and evoked levels of SP were calculated as a percentage of the total cellular content of SP per culture. Following field stimulation for 5 min, the amount of SP in the bathing solution was increased by ca. 15-fold over baseline.

Dihydropyridines modulate DRG cell calcium currents and neuropeptide secretion

Previous single-channel recording studies of chick DRG cells demonstrated that Bay K 8644, a DHP calcium-channel agonist, facilitates current flow through a specific subtype of calcium channel. These high-threshold, slowly inactivating calcium channels were designated as L-type by Nowycky et al. (1985a, b). In addition, Rane et al. (1987) recently reported that current flow through these same channels is inhibited by nifedipine, a DHP calcium-channel antagonist. We examined the effects of DHPs on DRG cell action potentials and calcium currents in order to determine whether their effects on calcium influx might be correlated with their effects on the release of SP.

Figure 6 illustrates the effects of DHPs on DRG cell action potentials and L-type calcium currents recorded from cells bathed in the same extracellular solution as was used in the release experiments. Intracellular microelectrode recordings demonstrated that Bay K 8644 (5 μ M) increased the duration of the

Figure 4. Effects of divalent cations on the electrically evoked release of SP. Nine cultures from the same plating were divided into 3 sets of 3, and each set was bathed in HBS containing either 3 mM Ca^{2+} , 1 mM Ba^{2+} plus 2 mM Ca^{2+} , or 0.1 mM Ca^{2+} plus 5 mM Co^{2+} , as indicated. Baseline (shaded columns) and evoked (unshaded columns) levels of SP were determined after 5 min exposure to each solution. Stimuli were delivered at 1 Hz. Each column indicates the mean \pm SEM for 3 cultures. The average cellular content of SP was $13,250 \pm 500$ pg/culture ($n = 9$). Inset, DRG cell action potentials recorded in each solution. Calibration bars, 10 msec, 20 mV. Note that Ba^{2+} increased the action potential duration and facilitated SP release, whereas Co^{2+} decreased the action potential duration and inhibited SP release.






Figure 5. Stimulus-response relationship for electrically evoked release of SP. Fifteen DRG cell cultures from the same plating were divided into 5 sets of 3, and each set was stimulated for either 60, 90, 150, 300, or 600 sec at 1 Hz. The HBS contained 1 mM Ba^{2+} and 2 mM Ca^{2+} . Each point indicates the mean \pm SEM value for 3 cultures. The average cellular content of SP was $12,800 \pm 600$ pg/culture ($n = 15$).

action potential in 14 of 18 cells tested, and increased the amplitude of the L-type calcium current (Fig. 6, A, B). In contrast, in 20 of 20 cells tested, nifedipine (5 μ M) failed to alter the duration of the action potential (Fig. 6C), but decreased the amplitude of the L-type calcium current in 9 of 13 cells (average decrease of $38 \pm 5\%$; mean \pm SEM) when the current was evoked from a relatively depolarized holding potential (-30 mV) that had been maintained for 5 sec or more, as illustrated in Figure 6E. In contrast, when the L-type current was evoked from a holding potential that approximated the normal resting potential of DRG neurons (-70 mV), nifedipine was relatively ineffective (Fig. 6D). These findings are consistent with the previously reported voltage and time-dependent actions of DHPs on L-type currents (Sanguinetti and Kass, 1984; Sanguinetti et al., 1986; Rane et al., 1987).

Next, we examined the effects of Bay K 8644 and nifedipine on SP release in order to determine whether the release process is triggered by calcium influx through L-type calcium channels. Neuropeptide secretion was induced by either field stimulation or depolarization with HBS containing 60 mM KCl. The amount of SP released was calculated as a percentage of the total cellular content of SP per culture prior to stimulation. As summarized in Table 1A, under conditions in which the HBS contained 2 mM $CaCl_2$ and 1 mM $BaCl_2$, treatment with Bay K 8644 (5 μ M) increased field stimulation-induced release of SP by 45% relative to control. In contrast, nifedipine (5 μ M, a concentration that is saturating for inhibition of L-type calcium currents in DRG neurons; Rane et al., 1987) did not significantly affect the release of SP, although an inhibitory trend was noted (Table 1A).

The failure of nifedipine to inhibit the electrically evoked release of SP can be explained by the fact that this DHP inhibits calcium-channel function in a voltage and time-dependent manner (cf. Fig. 6, D, E). That is, in previous studies of L-type calcium currents, the inhibitory action of nifedipine was accentuated by prior depolarization of the resting membrane potential (Sanguinetti and Kass, 1984; Rane et al., 1987). We tested for such a voltage and time-dependent action of nifedipine by examining its effects on the release of SP induced by prolonged depolarization of cultures with KCl.

When the release of SP was induced by 60 mM KCl, the actions of the DHPs became much more pronounced. In these exper-

iments, baseline values of SP were determined by exposing cultures for 5 min to HBS containing (in mM): 132 NaCl, 2.5 KCl, 2 $CaCl_2$, and 1 $BaCl_2$. For DHP-treated cultures, baseline values were determined in the presence of nifedipine or Bay K 8644. Next, cultures were depolarized for 5 min by HBS containing reduced NaCl (74 mM) and elevated KCl (60 mM). Intracellular recordings demonstrated that this concentration of KCl depolarized the resting membrane potential to -26 ± 1 mV ($n = 5$ neurons), thereby inducing the calcium-dependent release of SP. As summarized in Table 1B, Bay K 8644 (5 μ M) more than doubled the amount of SP released relative to control, whereas

Table 1. Effects of dihydropyridines on the release of substance P

Treatment	Baseline	Evoked	Change from control (%)
A. Field stimulation-induced release of SP			
Control ($n = 5$)	0.4 ± 0.1	9.6 ± 0.6	—
Nifedipine ($n = 5$)	0.6 ± 0.1	8.2 ± 0.9 (n.s.)	17% decrease
Bay K 8644 ($n = 5$)	0.4 ± 0.2	$13.7 \pm 0.8^*$	45% increase
B. 60 mM KCl-induced release of SP			
Control ($n = 5$)	0.5 ± 0.1	3.8 ± 0.5	—
Nifedipine ($n = 5$)	0.4 ± 0.2	$1.8 \pm 0.1^{**}$	58% decrease
Bay K 8644 ($n = 5$)	0.6 ± 0.2	$7.7 \pm 0.5^{**}$	115% increase

Results of 2 representative experiments using 30 cultures from 2 different platings. Each experiment assessed the effects of nifedipine and Bay K 8644 on the electrically and KCl-induced release of SP from cultures of the same plating. Values from both experiments were then averaged. Baseline and evoked levels of SP are expressed as a percentage of the total cellular content of SP per culture prior to stimulation. Values are the mean \pm SEM for the indicated number of cultures. A, Cultures were electrically stimulated for 5 min at 1 Hz, whereas B, cultures were depolarized for 5 min by HBS containing 60 mM KCl. The percentage change relative to control was calculated after subtracting baseline values from evoked values. Data were analyzed by Student's *t* test: * Value is significantly different from control: $p \leq 0.005$; ** $p \leq 0.001$; n.s., value is not significantly different from control: $p > 0.1$. The average cellular content was $14,200 \pm 250$ pg SP/culture ($n = 30$).




Figure 6. Effects of Bay K 8644 ($5 \mu\text{M}$) and nifedipine ($5 \mu\text{M}$) on DRG cell action potentials and L-type calcium currents. Bay K 8644 increased the action potential duration (*A*), whereas nifedipine was without effect (*C*). As shown in *B*, Bay K 8644 also increased the amplitude of the L-type calcium current (lower trace) when cells were held at -70 mV and depolarized to -15 mV for 40 msec (upper trace). *D, E*, Voltage and time-dependent inhibitory actions of nifedipine on L-type calcium currents. When cells were held at -70 mV and depolarized to 0 mV for 60 msec, nifedipine was a relatively weak antagonist (*D*). In contrast, when cells were held at -70 mV , stepped to -30 mV for 5 sec, and then depolarized to 0 mV for 60 msec, the inhibitory action of nifedipine was revealed (*E*). In *A–E* the bathing solution was HBS containing 2 mM Ca^{2+} and 1 mM Ba^{2+} . All examples are representative recordings obtained from different cells. Calibration bar: 50 mV , 40 msec (*A, C*); 4 nA , 25 msec (*B, D*); 0.8 nA , 25 msec (*E*).

nifedipine ($5 \mu\text{M}$) reduced the amount of SP released by 58%. At these concentrations, neither Bay K 8644 nor nifedipine significantly affected baseline levels of SP.

Conus toxin blocks DRG cell calcium currents and substance P release

It was recently reported that the molluscan neurotoxin $\omega\text{-CgTx}$, isolated from *Conus geographus* by Olivera et al. (1984), inhibits calcium influx associated with DRG action potentials (Kerr and Yoshikami, 1984) and blocks L- and N-type calcium currents in DRG cell bodies (McCleskey et al., 1986; Feldman et al., 1987). To further substantiate a role for L-type calcium channels in neuropeptide secretion, we examined the effects of $\omega\text{-CgTx}$ on the electrically evoked release of SP. As illustrated in Figure 7, *A–C*, $\omega\text{-CgTx}$ ($1 \mu\text{M}$) decreased the duration of DRG cell action potentials, irreversibly blocked the L-type calcium current, and inhibited the electrically evoked release of SP.

Discussion

Calcium-dependent release of substance P

The findings of the present study confirm previous reports that embryonic chick DRG neurons synthesize and secrete authentic SP when grown in primary dissociated cell culture (Mudge, 1979,




Figure 7. Analysis of the effects of $\omega\text{-CgTx}$. Application of $\omega\text{-CgTx}$ ($1 \mu\text{M}$) decreased the duration of the DRG cell action potential (*A*), decreased the amplitude of the L-type calcium current (*B*), and blocked the electrically evoked release of SP (*C*). Calcium currents were evoked from a holding potential of -70 mV in response to step depolarizations to 0 mV for 60 msec. In *A–C* the bathing solution was HBS containing 2 mM Ca^{2+} and 1 mM Ba^{2+} . For analysis of the effects of $\omega\text{-CgTx}$ on SP release, 6 cultures from the same plating were divided into 2 sets of 3. Each set was stimulated at 1 Hz for 3 min, and the amount of SP released was calculated as a percentage of the total cellular content of SP per culture prior to stimulation. Calibration bar: 80 mV , 40 msec (*A*); 0.8 nA , 25 msec (*B*).

1981; Mudge et al., 1979). These earlier studies examined the mechanism by which depolarizing concentrations of KCl induce neuropeptide secretion. We have extended these original observations by characterizing the electrically evoked release of SP in terms of its calcium-dependence, stimulus–response relationship, and its modulation by $\omega\text{-CgTx}$ and DHPs.

Three observations indicate that the electrically evoked release of SP does not result from nonspecific electropermeabilization of DRG cell membranes. First, intracellular recordings demonstrated that DRG neurons fire action potentials of normal configuration in response to electrical field stimulation. Secondly, the electrically evoked release of SP was fully repeatable, and, following stimulation, baseline levels of released SP returned to prestimulus levels, indicating that field stimulation does not damage the cells. Third, release was blocked by solutions containing either 5 mM Co^{2+} or $1 \mu\text{M } \omega\text{-CgTx}$. This is as expected for a process of calcium-dependent exocytosis, since little or no depolarization-induced calcium current is recorded from the soma membrane of DRG neurons bathed in such solutions (Dunlap and Fischbach, 1981; Fig. 7). Therefore, the

electrically evoked release of SP appears to be triggered by calcium influx through voltage-dependent calcium channels.

A role for L-type calcium channels in neuropeptide secretion

We have also demonstrated that the release of SP is facilitated by the DHP calcium-channel agonist Bay K 8644, and is inhibited by the DHP calcium-channel antagonist nifedipine. Significantly, it is now known that Bay K 8644 and nifedipine modulate slowly inactivating L-type calcium currents recorded from the soma membrane of DRG neurons (Nowycky et al., 1985a, b; Rane et al., 1987). In contrast, fast-inactivating (N- and T-type) calcium currents do not appear to be affected by either drug (Nowycky et al., 1985a, b; Rane et al., 1987). On the basis of these findings, we propose that DHPs modulate the release of SP by regulating calcium influx through L-type calcium channels located at release sites on primary afferent neurons.

It remains to be determined exactly where these release sites are located. For example, it might be argued that the release of SP is triggered by calcium influx through L-type calcium channels located in sensory nerve terminals. Alternatively, an action of Bay K 8644 or nifedipine on L-type calcium channels located in the somatic membrane of DRG neurons might explain the DHP sensitivity of the SP release mechanism. It is therefore important to recognize the limitations of release experiments conducted in dissociated cell culture where the sites of neuropeptide secretion remain undefined.

Voltage- and time-dependent actions of nifedipine

In the present study it was found that nifedipine inhibited the release of SP only under conditions of KCl-induced depolarization. No significant inhibitory action of the antagonist was observed when release was induced by electrical field stimulation. Moreover, nifedipine failed to inhibit calcium influx associated with DRG cell action potentials. These findings are explicable in terms of the voltage and time-dependent actions of DHPs on L-type calcium currents. Specifically, it was previously reported, and confirmed in the present study, that nifedipine inhibits L-type calcium currents most effectively when the membrane potential is held at *depolarized* levels (≤ -30 mV) for 5 sec or more (Sanguinetti and Kass, 1984; Rane et al., 1987). Clearly, such conditions are not achieved during electrical field stimulation, where SP release is triggered by action potentials of relatively short duration (10–50 msec) evoked from resting membrane potentials of -60 to -70 mV.

In contrast, when cultures were depolarized by 60 mM KCl, 5 μ M nifedipine inhibited the release of SP by 58%. Similar inhibitory actions of DHP calcium-channel antagonists were reported in previous studies examining KCl-induced calcium uptake into, or transmitter release from, rat DRG cells (Perney et al., 1986), cerebellar and retinal neurons (Takahashi and Ogura, 1983; Carboni et al., 1985), neuronal cell lines (Freedman et al., 1984), PC 12 cells (Toll, 1982), pituitary cells (Enyeart et al., 1985; Chang et al., 1986), and rat brain synaptosomes (Turner and Goldin, 1985).

Modulation of substance P release by Bay K 8644

The action of Bay K 8644 differed from that of nifedipine in that this calcium-channel agonist facilitated both the KCl-induced and electrically evoked release of SP. Furthermore, Bay K 8644 increased calcium influx associated with DRG cell action potentials. These findings are also explicable in terms of

the voltage and time-dependent actions of DHPs. Modulation of L-type calcium currents by Bay K 8644 does not require prior depolarization of the membrane potential. In fact, the agonist action of Bay K 8644 is most easily observed when L-type calcium currents are evoked from membrane potentials (-70 mV) close to the normal resting potential of DRG cells (see Fig. 6, A, B; see also Nowycky et al., 1985a, b; Sanguinetti et al., 1986). Such conditions are achieved during electrical field stimulation of DRG cells.

How these findings relate to previous studies examining the DHP sensitivity of neurosecretion

The findings reported herein are noteworthy in that they emphasize the limited usefulness of DHPs as pharmacological probes for L-type calcium channels. In particular, we have demonstrated that nifedipine inhibits, but does not completely block, the L-type calcium current, and that this antagonist action is both voltage and time-dependent. An understanding of such limitations is essential in order to properly evaluate previous studies reporting the failure of DHP calcium-channel antagonists to inhibit KCl-induced transmitter release from synaptosomal and brain slice preparations (reviewed by Miller, 1985). This failure to block secretion has been taken to indicate that in the central nervous system, calcium influx through L-type calcium channels does not normally trigger transmitter release (reviewed by Miller, 1987).

The findings of the present study suggest an alternative interpretation. We propose that the relatively slow antagonist action of DHPs (as reported here, and previously by Sanguinetti and Kass, 1984; Rane et al., 1987) severely limits their ability to inhibit transmitter release triggered by calcium influx during the initial phase (i.e., first 5 sec) of KCl-induced depolarization. This model predicts that antagonist DHPs will be effective only when the time course of transmitter release overlaps with the onset of L-type calcium-channel blockade, as is apparently the case during KCl-induced release of SP from DRG cell cultures.

In contrast, in those systems in which calcium influx and transmitter release are very transient in nature, no effect of antagonist DHPs is to be expected. For example, it was previously reported that DHPs fail to inhibit KCl-induced calcium uptake into rat brain synaptosomal preparations (Nachshen and Blaustein, 1979; Daniell et al., 1983; Reynolds et al., 1986; Suszkiw et al., 1986). However, it is also well documented that the rapid phase of KCl-induced calcium uptake into, and transmitter release from, synaptosomes is primarily limited to the first 1–2 sec of sustained depolarization (Drapeau and Blaustein, 1983; Floor, 1983; Turner and Goldin, 1985; Suszkiw et al., 1986). Our findings indicate that this time course of calcium influx may be too transient to allow effective blockade of the synaptosomal release mechanism by DHPs. In fact, such transient kinetics may result from rapid calcium-dependent inactivation of L-type calcium channels as intracellular levels of free calcium rise (Eckert and Chad, 1984). For these reasons, we wish to emphasize that in future studies examining the actions of DHPs, it will be necessary to obtain detailed information regarding the time course of calcium influx and transmitter release.

What is the role of N- and T-type calcium channels?

An important question that remains to be answered is whether calcium influx through N- or T-type calcium channels also triggers the release of SP from DRG neurons. On the basis of previous studies examining N- and T-type calcium currents in

the chick DRG soma membrane (Nowycky et al., 1985a), we would expect such a mechanism of release to exhibit DHP insensitivity. In fact, we found that 5 μ M nifedipine inhibited the KCl-induced release of SP by only 58%. Moreover, the electrically evoked release of SP was relatively insensitive to nifedipine. Although these observations are most simply explained by the voltage and time-dependent actions of DHPs on L-type calcium channels (as summarized above), it remains possible that DHP-insensitive (N- or T-type) calcium channels also play a role in the secretion process.

For example, it might be argued that the electrically evoked (and therefore physiologically relevant) release of SP results from calcium influx through N- or T-type channels, whereas L-type channels mediate secretion induced by KCl (see Miller, 1987). Our observation that ω -CgTx blocked the electrically evoked release of SP appears to rule out such a role for T-type channels, since current flow through this channel subtype was reported to be insensitive to the toxin (McCleskey et al., 1987). Therefore, the release of SP most likely results from Ca^{2+} influx through either N- or L-type channels, or possibly both. It must be emphasized, however, that the ω -CgTx effects reported herein do not resolve the question of which channel subtype is most important to the secretory process. Micromolar concentrations of ω -CgTx were previously reported to block both N- and L-type calcium currents (McCleskey et al., 1987). Therefore, this question will only be answered when more specific pharmacological probes for N- and L-type calcium channels are discovered.

Transmitter modulation of DHP-sensitive calcium channels

It was previously reported that norepinephrine (NE) and GABA inhibit DHP-sensitive, L-type calcium currents recorded from chick DRG cell bodies (Dunlap and Fischbach, 1981; Holz et al., 1986a; Rane et al., 1987). Since NE and GABA also inhibit the calcium-dependent release of SP from chick DRG cell cultures (Dunlap et al., 1986; Holz et al., 1986b), the transmitters may presynaptically inhibit neuropeptide secretion by regulating calcium influx through L-type calcium channels located at release sites on sensory nerve terminals (Dunlap et al., 1987). The findings of the present study support this concept. We have demonstrated that the depolarization-induced release of SP exhibits DHP sensitivity, as would be expected if calcium influx through L-type calcium channels triggers neuropeptide secretion. It may be concluded that in primary afferent neurons, the regulation of L-type calcium channels by neurotransmitters is likely to play an important role in processes governing the inhibition or facilitation of neurosecretion.

References

- Carboni, E., W. J. Wojcik, and E. Costa (1985) Dihydropyridines change the uptake of calcium induced by depolarization into primary cultures of cerebellar granule cells. *Neuropharmacology* 24: 1123–1126.
- Chang, M. M., and S. E. Leeman (1970) Isolation of a sialogogic peptide from bovine hypothalamic tissue and its characterization as substance P. *J. Biol. Chem.* 245: 4784–4790.
- Chang, J. P., E. M. McCoy, J. Graeter, K. Tasaka, and K. Catt (1986) Participation of voltage-dependent calcium channels in the action of gonadotropin-releasing hormone. *J. Biol. Chem.* 261: 9105–9108.
- Daniell, L. C., E. M. Barr, and S. W. Leslie (1983) Ca uptake into rat whole brain synaptosomes unaltered by dihydropyridine calcium antagonists. *J. Neurochem.* 41: 1455–1459.
- Dichter, M. A., and G. D. Fischbach (1977) The action potential of chick dorsal root ganglion neurones maintained in cell culture. *J. Physiol. (Lond.)* 267: 281–298.
- Drapeau, P., and M. P. Blaustein (1983) Initial release of dopamine from rat striatal synaptosomes: Correlation with calcium entry. *J. Neurosci.* 3: 703–713.
- Dunlap, K., and G. D. Fischbach (1978) Neurotransmitters decrease the calcium component of sensory neurone action potentials. *Nature* 276: 837–839.
- Dunlap, K., and G. D. Fischbach (1981) Neurotransmitters decrease the calcium conductance activated by depolarization of embryonic chick sensory neurons. *J. Physiol. (Lond.)* 317: 519–535.
- Dunlap, K., R. M. Kream, and G. G. Holz IV (1986) Alpha-2 adrenergic and GABA-B receptors mediate transmitter inhibition of neuropeptide secretion from dorsal root ganglion cells. *Soc. Neurosci. Abstr.* 12: 1195.
- Dunlap, K., S. G. Rane, and G. G. Holz IV (1987) Functional implications of calcium channel modulation in embryonic dorsal root ganglion neurons. In *Ion Channel Modulation*, A. D. Grinnell, D. L. Armstrong, and M. B. Jackson, eds., Plenum (in press).
- Eckert, R., and J. E. Chad (1984) Inactivation of Ca channels. *Prog. Biophys. Mol. Biol.* 44: 215–267.
- Enyeart, J. J., T. Aizawa, and P. M. Hinkle (1985) Dihydropyridine Ca antagonists: Potent inhibitors of secretion from normal and transformed pituitary cells. *Am. J. Physiol.* 248: C510–C519.
- Feldman, D. H., B. M. Olivera, and D. Yoshikami (1987) Omega conus toxin: A peptide that blocks calcium channels. *FEBS Lett.* 214: 295–300.
- Floor, E. (1983) Substance P release from K-depolarized rat brain synaptosomes at one second resolution. *Brain Res.* 279: 321–324.
- Freedman, S. B., G. Dawson, M. L. Villereal, and R. J. Miller (1984) Identification and characterization of voltage-sensitive calcium channels in neuronal clonal cell lines. *J. Neurosci.* 4: 1453–1467.
- Gamse, R., A. Molnar, and F. Lembeck (1979) Substance P release from spinal cord slices by capsaicin. *Life Sci.* 25: 629–636.
- Goetzl, E. J., T. Chernov, F. Renold, and D. G. Payan (1985) Neuropeptide regulation of the expression of immediate hypersensitivity. *J. Immunol.* 135: 802s–805s.
- Henry, J. L. (1977) Substance P and pain: A possible relation in afferent transmission. In *Substance P*, U.S. von Euler and B. Pernow, eds., pp. 231–239, Raven, New York.
- Henry, J. L. (1982) Relation of substance P to pain transmission: Neurophysiological evidence. *Ciba Found. Symp.* 91: 206–224.
- Holz, G. G., IV, S. G. Rane, and K. Dunlap (1986a) GTP-binding proteins mediate transmitter inhibition of voltage-dependent calcium channels. *Nature* 319: 670–672.
- Holz, G. G., IV, K. Dunlap, and R. M. Kream (1986b) Pertussis toxin-insensitive GTP-binding proteins couple alpha-2 and GABA-B receptors to inhibition of neurosecretion in dorsal root ganglion cells. *Soc. Neurosci. Abstr.* 12: 1195.
- Jessell, T. M., and L. L. Iversen (1977) Opiate analgesics inhibit substance P release from rat trigeminal nucleus. *Nature* 268: 549–551.
- Kerr, L. M., and D. Yoshikami (1984) A venom peptide with a novel presynaptic blocking action. *Nature* 308: 282–284.
- Kream, R. M., T. A. Schoenfeld, R. Mancuso, A. Clancy, W. El-Bermani, and F. Macrides (1985) Precursor forms of substance P (SP) in nervous tissue: Detection with antisera to SP, SP-Gly, and SP-Gly-Lys. *Proc. Natl. Acad. Sci. USA* 82: 4832–4836.
- McCleskey, E. W., A. P. Fox, D. Feldman, B. M. Olivera, R. W. Tsien, and D. Yoshikami (1987) ω -Conotoxin: Direct and persistent blockade of specific types of calcium channels in neurons but not muscle. *Proc. Natl. Acad. Sci. USA* 84: 4327–4331.
- Miller, R. J. (1985) How many types of calcium channels exist in neurons? *Trends Neurosci.* 80: 45–47.
- Miller, R. J. (1987) Multiple calcium channels and neuronal function. *Science* 235: 46–52.
- Mudge, A. W. (1979) Studies on substance P, somatostatin, and enkephalin in cultures of sensory neurons. Ph.D. dissertation thesis, Harvard University, Cambridge, MA.
- Mudge, A. W. (1981) Effect of chemical environment on levels of substance P and somatostatin in cultured sensory neurones. *Nature* 292: 764–767.
- Mudge, A. W., S. E. Leeman, and G. D. Fischbach (1979) Enkephalin inhibits release of substance P from sensory neurons in culture and decreases action potential duration. *Proc. Natl. Acad. Sci. USA* 76: 526–530.
- Nachshen, D. A., and M. P. Blaustein (1979) The effects of some organic “calcium antagonists” on calcium influx in presynaptic nerve terminals. *Mol. Pharmacol.* 16: 579–586.

- Nicoll, R. A., C. Schenker, and S. E. Leeman (1980) Substance P as a transmitter candidate. *Annu. Rev. Neurosci.* 3: 227-268.
- Nowycky, M. C., A. P. Fox, and R. W. Tsien (1985a) Three types of neuronal calcium channel with different calcium agonist sensitivity. *Nature* 316: 440-443.
- Nowycky, M. C., A. P. Fox, and R. W. Tsien (1985b) Long-opening mode of gating of neuronal calcium channels and its promotion by the dihydropyridine calcium agonist Bay K 8644. *Proc. Natl. Acad. Sci. USA* 82: 2178-2182.
- Olivera, B. M., J. M. McIntosh, L. J., Cruz, F. A. Luque, and W. R. Gray (1984) Purification and sequence of a presynaptic peptide toxin from *Conus geographus* venom. *Biochemistry* 23: 5087-5090.
- Otsuka, M., and S. Konishi (1976) Release of substance P-like immunoreactivity from isolated spinal cord of newborn rat. *Nature* 264: 83-84.
- Payan, D. G., and E. J. Goetzl (1985) Modulation of lymphocyte function by sensory neuropeptides. *J. Immunol.* 135: 783s-786s.
- Perney, T. M., L. D. Hirning, and R. J. Miller (1986) Multiple calcium channels mediate neurotransmitter release from peripheral neurons. *Proc. Natl. Acad. Sci. USA* 83: 6656-6659.
- Pernow, B. (1983a) Substance P. *Pharmacol. Rev.* 35: 85-141.
- Pernow, B. (1983b) Substance P—a putative mediator of antidromic vasodilation. *Gen. Pharmacol.* 14: 13-16.
- Pernow, B. (1985) Role of tachykinins in neurogenic inflammation. *J. Immunol.* 135: 812s-815s.
- Rane, S. G., and K. Dunlap (1986) Kinase C activator 1,2-oleoyl-acetyl-glycerol attenuates voltage-dependent calcium current in sensory neurons. *Proc. Natl. Acad. Sci. USA* 83: 184-188.
- Rane, S. G., G. G. Holz IV, and K. Dunlap (1987) Dihydropyridine inhibition of neuronal calcium current and substance P release. *Pflügers Arch.* 409: 361-366.
- Reynolds, I. J., J. A. Wagner, S. H. Snyder, S. A. Thayer, B. M. Olivera, and R. J. Miller (1986) Brain voltage-sensitive calcium channel subtypes differentiated by ω -conotoxin fraction GVIA. *Proc. Natl. Acad. Sci. USA* 83: 8804-8807.
- Sanguinetti, M. C., and R. S. Kass (1984) Voltage-dependent block of calcium channel current in the calf cardiac Purkinje fiber by dihydropyridine calcium channel antagonists. *Circ. Res.* 55: 336-348.
- Sanguinetti, M. C., D. S. Krafte, and R. S. Kass (1986) Voltage-dependent modulation of Ca current in heart cells by Bay K 8644. *J. Gen. Physiol.* 88: 369-392.
- Suszkiew, J. B., M. E. O'Leary, M. M. Murawsky, and T. Wang (1986) Presynaptic calcium channels in rat cortical synaptosomes: Fast-kinetics of phasic calcium influx, channel inactivation, and relationship to nitrendipine receptors. *J. Neurosci.* 6: 1349-1357.
- Takahashi, M., and A. Ogura (1983) Dihydropyridines as potent calcium channel blockers in neuronal cells. *FEBS Lett.* 152: 191-194.
- Toll, L. (1982) High-affinity binding and inhibition of calcium transport in clonal cell line. *J. Biol. Chem.* 257: 13189-13192.
- Turner, T. J., and S. M. Goldin (1985) Calcium channels in rat brain synaptosomes: Identification and pharmacological characterization: High affinity blockade by organic Ca channel blockers. *J. Neurosci.* 5: 841-895.
- Yaksh, T. L., T. M. Jessell, R. Gamse, A. W. Mudge, and S. E. Leeman (1980) Intrathecal morphine inhibits substance P release from mammalian spinal cord *in vivo*. *Nature* 286: 155-157.